



# An optimal double inequality between geometric and identric means

Miao-Kun Wang<sup>a</sup>, Zi-Kui Wang<sup>b</sup>, Yu-Ming Chu<sup>a,\*</sup>

<sup>a</sup> Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China

<sup>b</sup> Department of Mathematics, Hangzhou Normal University, Hangzhou 310018, China

## ARTICLE INFO

### Article history:

Received 6 June 2011

Received in revised form 10 September 2011

Accepted 19 September 2011

### Keywords:

Geometric mean

Identric mean

Inequality

## ABSTRACT

We find the greatest value  $p$  and least value  $q$  in  $(0, 1/2)$  such that the double inequality  $G(pa + (1-p)b, pb + (1-p)a) < I(a, b) < G(qa + (1-q)b, qb + (1-q)a)$  holds for all  $a, b > 0$  with  $a \neq b$ . Here,  $G(a, b)$ , and  $I(a, b)$  denote the geometric, and identric means of two positive numbers  $a$  and  $b$ , respectively.

© 2011 Elsevier Ltd. All rights reserved.

## 1. Introduction

The classical geometric mean  $G(a, b)$  and identric mean  $I(a, b)$  of two positive numbers  $a$  and  $b$  with  $a \neq b$  are defined by

$$G(a, b) = \sqrt{ab} \quad \text{and} \quad I(a, b) = \frac{1}{e} \left( \frac{b^b}{a^a} \right)^{1/(b-a)},$$

respectively. Recently, both mean values have been the subject of intensive research. In particular, many remarkable inequalities for  $G$  and  $I$  can be found in the literature [1–21].

Let  $M_p(a, b) = [(a^p + b^p)/2]^{1/p}$  ( $p \neq 0$ ) and  $M_0(a, b) = \sqrt{ab}$ ,  $H(a, b) = 2ab/(a + b)$ ,  $A(a, b) = (a + b)/2$ ,  $L(a, b) = (a - b)/(\log a - \log b)$ , and  $P(a, b) = (a - b)/(4 \arctan \sqrt{a/b} - \pi)$  be the  $p$ -th power, harmonic, arithmetic, logarithmic, and Seiffert means of two positive numbers  $a$  and  $b$  with  $a \neq b$ , respectively. Then it is well known that

$$\min\{a, b\} < H(a, b) = M_{-1}(a, b) < G(a, b) = M_0(a, b) < L(a, b) \\ < P(a, b) < I(a, b) < A(a, b) = M_1(a, b) < \max\{a, b\}$$

for all  $a, b > 0$  with  $a \neq b$ .

The following sharp bounds for  $I$ ,  $(LI)^{1/2}$ , and  $(L + I)/2$  in terms of power mean are presented in [22].

$$M_{2/3}(a, b) < I(a, b) < M_{\log 2}(a, b), \quad M_0(a, b) < \sqrt{L(a, b)I(a, b)} < M_{1/2}(a, b), \\ M_{\log 2/(1+\log 2)}(a, b) < [L(a, b) + I(a, b)]/2 < M_{1/2}(a, b)$$

for all  $a, b > 0$  with  $a \neq b$ .

In [23], Alzer and Qiu proved that the inequalities

$$\alpha A(a, b) + (1 - \alpha)G(a, b) < I(a, b) < \beta A(a, b) + (1 - \beta)G(a, b) \quad (1.1)$$

hold for all positive real numbers  $a$  and  $b$  with  $a \neq b$  if and only if  $\alpha \leq 2/3$  and  $\beta \geq 2/e = 0.73575 \dots$ .

\* Corresponding author.

E-mail address: [chuyuming@hutczj.cn](mailto:chuyuming@hutczj.cn) (Y.-M. Chu).

In [24–26], the authors answered the questions: for  $\alpha, \beta \in (0, 1)$  with  $\alpha + \beta < 1$ , what are the greatest values  $p_1, p_2, p_3$  and the least values  $q_1, q_2, q_3$  such that  $M_{p_1}(a, b) < P^\alpha(a, b)G^{1-\alpha}(a, b) < M_{q_1}(a, b), M_{p_2}(a, b) < G^\alpha(a, b)L^{1-\alpha}(a, b) < M_{q_2}(a, b)$ , and  $M_{p_3}(a, b) < A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b) < M_{q_3}(a, b)$  for all  $a, b > 0$  with  $a \neq b$ ?

For fixed  $a, b > 0$  with  $a \neq b$  and  $x \in [0, 1/2]$ , let

$$g(x) = G(xa + (1-x)b, xb + (1-x)a).$$

Then it is not difficult to verify that  $g(x)$  is continuous and strictly increasing in  $[0, 1/2]$ . Note that  $g(0) = G(a, b) < I(a, b)$  and  $g(1/2) = A(a, b) > I(a, b)$ . Therefore, it is natural to ask what are the greatest value  $p$  and least value  $q$  in  $(0, 1/2)$  such that the double inequality  $G(pa + (1-p)b, pb + (1-p)a) < I(a, b) < G(qa + (1-q)b, qb + (1-q)a)$  holds for all  $a, b > 0$  with  $a \neq b$ . The main purpose of this paper is to answer this question. Our main result is the following Theorem 1.1.

**Theorem 1.1.** *If  $p, q \in (0, 1/2)$ , then the double inequality*

$$G(pa + (1-p)b, pb + (1-p)a) < I(a, b) < G(qa + (1-q)b, qb + (1-q)a) \quad (1.2)$$

*holds for all  $a, b > 0$  with  $a \neq b$  if and only if  $p \leq (1 - \sqrt{1 - 4/e^2})/2$  and  $q \geq (3 - \sqrt{3})/6$ .*

## 2. Proof of Theorem 1.1

**Proof of Theorem 1.1.** Let  $\lambda = (3 - \sqrt{3})/6$  and  $\mu = (1 - \sqrt{1 - 4/e^2})/2$ . Then from the monotonicity of the function  $g(x) = G(xa + (1-x)b, xb + (1-x)a)$  in  $[0, 1/2]$  we know that to prove inequality (1.2) we only need to prove that inequalities

$$I(a, b) < G(\lambda a + (1-\lambda)b, \lambda b + (1-\lambda)a) \quad (2.1)$$

and

$$I(a, b) > G(\mu a + (1-\mu)b, \mu b + (1-\mu)a) \quad (2.2)$$

hold for all  $a, b > 0$  with  $a \neq b$ .

Without loss of generality, we assume that  $a > b$ . Let  $t = a/b > 1$  and  $r \in (0, 1/2)$ , then simple computation leads to

$$\begin{aligned} & \log G(ra + (1-r)b, rb + (1-r)a) - \log I(a, b) \\ &= \frac{1}{2} \log \left( \frac{ra}{b} + 1 - r \right) + \frac{1}{2} \log \left[ \frac{(1-r)a}{b} + r \right] + \log b - \frac{a(\log a - \log b)}{a-b} - \log b + 1 \\ &= \frac{1}{2} \log(rt + 1 - r) + \frac{1}{2} \log[(1-r)t + r] - \frac{t}{t-1} \log t + 1. \end{aligned} \quad (2.3)$$

Let

$$f(t) = \frac{1}{2} \log(rt + 1 - r) + \frac{1}{2} \log[(1-r)t + r] - \frac{t}{t-1} \log t + 1, \quad (2.4)$$

then

$$f(1) = 0, \quad (2.5)$$

$$\lim_{t \rightarrow +\infty} f(t) = \frac{1}{2} \log[r(1-r)] + 1, \quad (2.6)$$

$$f'(t) = \frac{f_1(t)}{2(t-1)^2}, \quad (2.7)$$

where

$$\begin{aligned} f_1(t) &= 2 \log t - \frac{t^2 - 1}{[rt + (1-r)][(1-r)t + r]}, \\ f_1(1) &= 0, \end{aligned} \quad (2.8)$$

$$\lim_{t \rightarrow +\infty} f_1(t) = +\infty, \quad (2.9)$$

$$f_1'(t) = \frac{f_2(t)}{t[rt + (1-r)]^2[(1-r)t + r]^2}, \quad (2.10)$$

where

$$f_2(t) = 2r^2(1-r)^2t^4 - (2r^2 - 2r + 1)(1-2r)^2t^3 + 2(6r^4 - 12r^3 + 12r^2 - 6r + 1)t^2 - (2r^2 - 2r + 1)(1-2r)^2t + 2r^2(1-r)^2,$$

$$f_2(1) = 0,$$

$$\lim_{t \rightarrow +\infty} f_2(t) = +\infty, \quad (2.11)$$

$$\lim_{t \rightarrow +\infty} f_2(t) = +\infty, \quad (2.12)$$

$$f_2'(t) = 8r^2(1-r)^2t^3 - 3(2r^2 - 2r + 1)(1-2r)^2t^2 + 4(6r^4 - 12r^3 + 12r^2 - 6r + 1)t - (2r^2 - 2r + 1)(1-2r)^2,$$

$$f_2'(1) = 0, \quad \lim_{t \rightarrow +\infty} f_2'(t) = +\infty, \quad (2.13)$$

$$f_2''(t) = 24r^2(1-r)^2t^2 - 6(2r^2 - 2r + 1)(1-2r)^2t + 4(6r^4 - 12r^3 + 12r^2 - 6r + 1),$$

$$f_2''(1) = -2(6r^2 - 6r + 1), \quad (2.14)$$

$$\lim_{t \rightarrow +\infty} f_2''(t) = +\infty, \quad (2.15)$$

$$f_2'''(t) = 48r^2(1-r)^2t - 6(2r^2 - 2r + 1)(1-2r)^2, \quad (2.16)$$

$$f_2'''(1) = -6(6r^2 - 6r + 1) \quad (2.17)$$

and

$$\lim_{t \rightarrow +\infty} f_2'''(t) = +\infty. \quad (2.18)$$

We divide the proof into two cases.

Case 1.  $r = \lambda = (3 - \sqrt{3})/6$ . Then Eqs. (2.14) and (2.17) lead to

$$f_2''(1) = 0 \quad (2.19)$$

and

$$f_2'''(1) = 0. \quad (2.20)$$

It follows from (2.16) that  $f_2'''(t)$  is strictly increasing in  $[1, +\infty)$ , then (2.20) implies that  $f_2'''(t) > 0$  for  $t \in (1, +\infty)$ . Hence,  $f_2''(t)$  is strictly increasing in  $[1, +\infty)$ .

From Eqs. (2.5), (2.7), (2.8), (2.10), (2.11), (2.13) and (2.19) together with the monotonicity of  $f_2''(t)$  we clearly see that

$$f(t) > 0 \quad (2.21)$$

for  $t \in (1, +\infty)$ .

Therefore, inequality (2.1) follows from Eqs. (2.3) and (2.4) together with inequality (2.21).

Case 2.  $r = \mu = (1 - \sqrt{1 - 4/e^2})/2$ . Then Eqs. (2.14) and (2.17) lead to

$$f_2''(1) = 2\left(\frac{6}{e^2} - 1\right) < 0 \quad (2.22)$$

and

$$f_2'''(1) = 6\left(\frac{6}{e^2} - 1\right) < 0. \quad (2.23)$$

It follows from (2.18) and (2.23) together with the monotonicity of  $f_2'''(t)$  that there exists  $t_1 > 1$  such that  $f_2'''(t) < 0$  for  $t \in (1, t_1)$  and  $f_2'''(t) > 0$  for  $t \in (t_1, +\infty)$ . Hence,  $f_2''(t)$  is strictly decreasing in  $[1, t_1]$  and strictly increasing in  $[t_1, +\infty)$ .

From (2.15) and (2.22) together with the piecewise monotonicity of  $f_2''(t)$  we clearly see that there exists  $t_2 > t_1 > 1$  such that  $f_2'(t)$  is strictly decreasing in  $[1, t_2]$  and strictly increasing in  $[t_2, +\infty)$ . Then (2.13) leads to the conclusion that there exists  $t_3 > t_2 > 1$  such that  $f_2(t)$  is strictly decreasing in  $[1, t_3]$  and strictly increasing in  $[t_3, +\infty)$ .

It follows from (2.10) to (2.12) and the piecewise monotonicity of  $f_2(t)$  that there exists  $t_4 > t_3 > 1$  such that  $f_1(t)$  is strictly decreasing in  $[1, t_4]$  and strictly increasing in  $[t_4, +\infty)$ .

From (2.7) to (2.9) and the piecewise monotonicity of  $f_1(t)$  we know that there exists  $t_5 > t_4 > 1$  such that  $f(t)$  is strictly decreasing in  $[1, t_5]$  and strictly increasing in  $[t_5, +\infty)$ .

Note that Eq. (2.6) becomes

$$\lim_{t \rightarrow +\infty} f(t) = 0 \quad (2.24)$$

for  $r = \mu = (1 - \sqrt{1 - 4/e^2})/2$ .

From (2.5) and (2.24) together with the piecewise monotonicity of  $f(t)$  we conclude that

$$f(t) < 0 \quad (2.25)$$

for  $t > 1$ .

Therefore, inequality (2.2) follows from Eqs. (2.3) and (2.4) together with inequality (2.25).

Next, we prove that  $\lambda = (3 - \sqrt{3})/6$  is the best possible parameter in  $(0, 1/2)$  such that inequality (2.1) holds for all  $a, b > 0$  with  $a \neq b$ . In fact, if  $0 < r < \lambda = (3 - \sqrt{3})/6$ , then (2.14) leads to  $f_2''(1) = -2(6r^2 - 6r + 1) < 0$ . Thus from the continuity of  $f_2''(t)$  we know that there exists  $\delta > 0$  such that

$$f_2''(t) < 0 \quad (2.26)$$

for  $t \in (1, 1 + \delta)$ .

It follows from (2.3) to (2.5), (2.7), (2.8), (2.10), (2.11), (2.13) and (2.26) that  $I(a, b) > G(ra + (1 - r)b, rb + (1 - r)a)$  for  $a/b \in (1, 1 + \delta)$ .

Finally, we prove that the parameter  $\mu = (1 - \sqrt{1 - 4/e^2})/2$  is the best possible parameter in  $(0, 1/2)$  such that inequality (2.2) holds for all  $a, b > 0$  with  $a \neq b$ . In fact, if  $(1 - \sqrt{1 - 4/e^2})/2 = \mu < r < 1/2$ , then (2.6) leads to the conclusion that  $\lim_{t \rightarrow +\infty} f(t) > 0$ . Hence there exists  $T > 1$  such that

$$f(t) > 0 \quad (2.27)$$

for  $t \in (T, +\infty)$ .

Therefore,  $I(a, b) < G(ra + (1 - r)b, rb + (1 - r)a)$  for  $a/b \in (T, +\infty)$  follows from Eqs. (2.3) and (2.4) together with inequality (2.27).  $\square$

**Remark 2.1.** The lower and upper bounds in (1.2) with  $p = (1 - \sqrt{1 - 4/e^2})/2$  and  $q = (3 - \sqrt{3})/6$  are better than those in (1.1) with  $\alpha = 2/3$  and  $\beta = 2/e$  for some  $a, b > 0$ , respectively. In fact, if we let  $t = \sqrt{a/b} > 0$ , then

$$\begin{aligned} & [G(pa + (1 - p)b, pb + (1 - p)a)]^2 - [\alpha A(a, b) + (1 - \alpha)G(a, b)]^2 \\ &= [pa + (1 - p)b][pb + (1 - p)a] - \left[ \frac{a + b}{3} + \frac{\sqrt{ab}}{3} \right]^2 \\ &= \frac{b^2(t - 1)^2}{9e^2} [(9 - e^2)t^2 + 2(9 - 2e^2)t + 9 - e^2] \\ &= \frac{b^2(9 - e^2)(t - 1)^2}{9e^2} \left[ t - \frac{2e^2 - 9 - \sqrt{3e^2(e^2 - 6)}}{9 - e^2} \right] \left[ t - \frac{2e^2 - 9 + \sqrt{3e^2(e^2 - 6)}}{9 - e^2} \right] \end{aligned} \quad (2.28)$$

and

$$\begin{aligned} & [G(qa + (1 - q)b, qb + (1 - q)a)]^2 - [\beta A(a, b) + (1 - \beta)G(a, b)]^2 \\ &= [qa + (1 - q)b][qb + (1 - q)a] - \left[ \frac{a + b}{e} + \frac{(e - 2)\sqrt{ab}}{e} \right]^2 \\ &= \frac{b^2(t - 1)^2}{6e^2} [(e^2 - 6)t^2 + 2(e^2 - 6e + 6)t + e^2 - 6] \\ &= \frac{b^2(e^2 - 6)(t - 1)^2}{6e^2} \left[ t - \frac{-e^2 + 6e - 6 - \sqrt{12e(e - 2)(3 - e)}}{e^2 - 6} \right] \\ &\quad \times \left[ t - \frac{-e^2 + 6e - 6 + \sqrt{12e(e - 2)(3 - e)}}{e^2 - 6} \right]. \end{aligned} \quad (2.29)$$

Eqs. (2.28) and (2.29) imply that  $G(pa + (1 - p)b, pb + (1 - p)a) > 2A(a, b)/3 + G(a, b)/3$  if and only if  $a/b < [2e^2 - 9 - \sqrt{3e^2(e^2 - 6)}]^2 / (9 - e^2)^2 = 0.020 \dots$  or  $a/b > [2e^2 - 9 + \sqrt{3e^2(e^2 - 6)}]^2 / (9 - e^2)^2 = 49.439 \dots$ , and  $G(qa + (1 - q)b, qb + (1 - q)a) < 2A(a, b)/e + (e - 2)G(a, b)/e$  if and only if  $[-e^2 + 6e - 6 - \sqrt{12e(e - 2)(3 - e)}]^2 / (e^2 - 6)^2 = 0.064 \dots < a/b < [-e^2 + 6e - 6 + \sqrt{12e(e - 2)(3 - e)}]^2 / (e^2 - 6)^2 = 15.619 \dots$ .  $\square$

## Acknowledgments

The authors wish to thank the anonymous referees for their careful reading of the manuscript and their fruitful comments and suggestions. This research was supported by the Natural Science Foundation of China under Grant 11071069 and the Innovation Team Foundation of the Department of Education of Zhejiang Province under Grant T200924.

## References

- [1] Y.-M. Chu, W.-F. Xia, Two sharp inequalities for power mean, geometric mean, and harmonic mean, *J. Inequal. Appl.* (2009) 6. Article ID 741923.
- [2] Y.-M. Chu, W.-F. Xia, Inequalities for generalized logarithmic means, *J. Inequal. Appl.* (2009) 7. Article ID 763252.
- [3] B.-Y. Long, Y.-M. Chu, Optimal inequalities for generalized logarithmic, arithmetic, and geometric means, *J. Inequal. Appl.* (2010) 10. Article ID 806825.
- [4] Y.-M. Chu, B.-Y. Long, Best possible inequalities between generalized logarithmic mean and classical means, *Abstr. Appl. Anal.* (2010) 13. Article ID 303286.
- [5] Y.-M. Chu, B.-Y. Long, Sharp inequalities between means, *Math. Inequal. Appl.* 14 (3) (2011) 647–655.
- [6] Y.-M. Chu, C. Zong, G.D. Wang, Optimal convex combination bounds of Seiffert and geometric means for the arithmetic mean, *J. Math. Inequal.* 5 (3) (2011) 429–434.
- [7] H. Alzer, Inequalities for arithmetic, geometric and harmonic means, *Bull. Lond. Math. Soc.* 22 (4) (1990) 362–366.
- [8] H.-J. Seiffert, Eine Integralungleichung für streng monotone Funktionen mit logarithmisch konvexer Umkehrfunktion, *Elem. Math.* 44 (1) (1989) 16–18.
- [9] C.-P. Chen, F. Qi, Monotonicity properties and inequalities of functions related to means, *Rocky Mountain J. Math.* 36 (3) (2006) 857–865.
- [10] L. Zhu, Some new inequalities for means in two variables, *Math. Inequal. Appl.* 11 (3) (2008) 443–448.
- [11] E. Neuman, J. Sándor, Companion inequalities for certain bivariate means, *Appl. Anal. Discrete Math.* 3 (1) (2009) 46–51.
- [12] H.-X. Du, Some inequalities for bivariate means, *Commun. Korean Math. Soc.* 24 (4) (2009) 553–559.
- [13] O. Kouba, New bounds for the identric mean of two arguments, *JIPAM. J. Inequal. Pure Appl. Math.* 9 (3) (2008) 6. Article 71.
- [14] T. Trif, On certain inequalities involving the identric mean in  $n$  variables, *Studia Univ. Babeş-Bolyai Math.* 46 (4) (2001) 105–114.
- [15] M.K. Vamanamurthy, M. Vuorinen, Inequalities for means, *J. Math. Anal. Appl.* 183 (1) (1994) 155–166.
- [16] J. Sándor, On the identric and logarithmic means, *Aequationes Math.* 40 (2–3) (1990) 261–270.
- [17] J. Sándor, A note on some inequalities for means, *Arch. Math.* 56 (5) (1991) 471–473.
- [18] J. Sándor, On refinements of certain inequalities for means, *Arch. Math. Brno* 31 (4) (1995) 279–282.
- [19] J. Sándor, Two inequalities for means, *Internat. J. Math. Math. Sci.* 18 (3) (1995) 621–623.
- [20] H. Alzer, Ungleichungen für Mittelwerte, *Arch. Math.* 47 (5) (1986) 422–426.
- [21] Y.-F. Qiu, M.-K. Wang, Y.-M. Chu, G.-D. Wang, Two sharp inequalities for Lehmer mean, identric mean and logarithmic mean, *J. Math. Inequal.* 5 (3) (2011) 301–306.
- [22] H. Alzer, Bestmögliche Abschätzungen für spezielle Mittelwerte, *Zb. Rad. Prir.-Mat. Fak. Ser. Mat.* 23 (1) (1993) 331–346.
- [23] H. Alzer, S.-L. Qiu, Inequalities for mean in two variables, *Arch. Math.* 80 (2) (2003) 201–215.
- [24] Y.-M. Chu, Y.-F. Qiu, M.-K. Wang, Sharp power mean bounds for the combination of Seiffert and geometric means, *Abstr. Appl. Anal.* (2010) 12. Article ID 108920.
- [25] M.-Y. Shi, Y.-M. Chu, Y.-P. Jiang, Optimal inequalities among various means of two arguments, *Abstr. Appl. Anal.* (2009) 10. Article ID 694394.
- [26] B.-Y. Long, Y.-M. Chu, Optimal power mean bounds for the weighted geometric mean of classical means, *J. Inequal. Appl.* (2010) 6. Article ID 905679.